

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

1012825

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

LITERATUUR KOPIEEN



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 31/335, 31/34, 31/27, 31/13, 31/12, 31/16	A1	(11) International Publication Number: WO 95/27485 (43) International Publication Date: 19 October 1995 (19.10.95)
(21) International Application Number: PCT/US95/02872 (22) International Filing Date: 6 March 1995 (06.03.95) (30) Priority Data: 08/224,949 8 April 1994 (08.04.94) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: BISSETT, Donald, L.; 3925 Dust Commander Drive, Hamilton, OH 45011 (US). SHAYA, Steven, Alan; 3618 Fawnrun Drive, Cincinnati, OH 45241 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHODS OF USING IRON CHELATING COMPOUNDS TO REDUCE FREE RADICAL DAMAGE IN MAMMALS (57) Abstract The subject invention involves a method of reducing free radical damage in mammals comprising administering to a mammal a composition comprising a safe and effective amount of a compound selected from 1-phenyl-1,2-propanedione-2-oxime, benzoylacetone, piroctone, 2-furildioxime, 2-furilmonoxime, diethyldithiocarbamic acid, deferoxamine and 1,2-dimethyl-3-hydroxy-pyrid-4-one, or a pharmaceutically acceptable salt thereof, or mixtures of the subject compounds and/or their salts.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

METHODS OF USING IRON CHELATING COMPOUNDS TO REDUCE FREE RADICAL DAMAGE IN MAMMALS

5

The subject invention relates to the field of protecting mammals from free radical damage and conditions accelerated by free radical damage. Specifically, the subject invention relates to novel methods of using certain iron-chelating compounds for reducing the level of free radicals in mammalian cells.

BACKGROUND OF THE INVENTION

Free radicals, in particular oxygen radicals, in mammalian cells arise from a variety of environmental sources. Such sources include smoke, pollution and radiation in addition to normal cell metabolism and inflammatory processes. Free radicals are known to be damaging to biological tissue. In the body, the potential for damage by free radicals is greatly increased by the presence of iron (and to a lesser extent by the presence of copper), which catalyzes their conversion to less stable and therefore more reactive radical species. The resulting effects of such radicals on components such as structural proteins, membrane lipids and nucleic acids include alteration or loss of tissue and cell function, cell death, and cancer. Based on a growing body of evidence, it is also believed that the accumulated damage from free radicals, including those produced as by-products in normal metabolism, is responsible for chronological aging. This has been termed the free radical theory of aging.

In vitro and in vivo studies on humans and other mammals also indicate that free radicals play an important role in the activation of certain human immunodeficiency virus (HIV) types. See, for example, Schreck, R., B. Meier, D. N. Manner, W. Dröge and P.A. Baeuerle, "Dithiocarbamates as Potent Inhibitors of Nuclear Factor κ B Activation in Intact Cells," J. Exp. Med., Vol. 175, pp. 1181-1194, (1992). Certain iron chelators have therefore been considered for the treatment of Acquired Immune Deficiency Syndrome (AIDS). See, for example, Hersh, E. M. et al., "Ditiocarb Sodium (Diethyldithiocarbamate) Therapy in Patients with Symptomatic HIV Infection and Aids," J. A. M. A., Vol. 265, pp.1538-1544, (1991).

To combat the damaging effects of free radicals, especially oxygen radicals, free radical scavengers and anti-oxidants have been used. These compounds react with the radical species to convert them to stable, nonreactive materials.

It is an object of the subject invention to provide methods for reducing free radical damage in mammalian cells.

It is also an object of the subject invention to provide compositions which reduce free radical damage in mammalian cells.

SUMMARY OF THE INVENTION

The subject invention involves a method of reducing free radical damage in mammals comprising administering to a mammal a composition comprising a safe and effective amount of a compound selected from 1-phenyl-1,2-propanedione-2-oxime, benzoylacetone, 5 piroctone, 2-furildioxime, 2-furilmonoxime, diethyldithiocarbamic acid, deferoxamine and 1,2-dimethyl-3-hydroxy-pyrid-4-one, or a pharmaceutically acceptable salt thereof, or mixtures of the subject compounds and/or their salts.

DETAILED DESCRIPTION OF THE INVENTION

10 It has been unexpectedly found that compositions comprising the subject compounds exhibit the ability to reduce levels of free radicals in mammals. While the subject invention is not limited to any particular mode of action, it is believed that the subject compounds may reduce the level of free radicals in mammalian cells by preventing the formation of the most damaging radical species. The subject compounds are believed to bind to iron such that iron 15 cannot participate in the formation of the above mentioned radical species. The subject compounds, which act as iron chelators, are unexpectedly effective against both environmentally induced radical production as well as the endogenous radical production arising from metabolism.

As used herein, "alkyl" means a substituted or unsubstituted carbon-containing chain 20 which may be straight or branched; saturated, monounsaturated (i.e., one double or triple bond in the chain), or polyunsaturated (i.e., two or more double bonds in the chain; two or more triple bonds in the chain; one or more double and one or more triple bonds in the chain).

As used herein, "topical application" means directly laying on or spreading on outer 25 skin.

As used herein, "pharmaceutically-acceptable" means that salts, drugs, medicaments or inert ingredients which the term describes are suitable for use in contact with the tissues of humans and lower animals without undue toxicity, incompatibility, instability, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio.

30 As used herein, "safe and effective amount" means an amount of compound or composition sufficient to significantly induce a positive modification in the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. The safe and effective amount of the compound or composition will vary with the particular condition being treated, the age and 35 physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the specific compound or composition employed, the particular pharmaceutically-acceptable carrier utilized, and like factors within the knowledge and expertise of the attending physician.

As used herein, "free radical" means an atom or group of atoms that have one or more unpaired electrons. Such atoms or groups of atoms are highly reactive and unstable species. Also included among "free radicals" are other oxygen species which are highly reactive toward biological systems. Some "free radicals" are produced by the catalytic action of metals (e.g., iron or copper) or are more reactive toward biological systems in the presence of metals (e.g., iron or copper). Specific non-limiting examples of "free radicals" as defined above are:

	superoxide	O_2^-
	hydroperoxyl radical	$HO_2\cdot$
10	peroxide ion	O_2^{2-}
	hydroperoxyl anion	HO_2^-
	hydroxyl radical	$HO\cdot$
	singlet oxygen	1O_2
	hydrogen peroxide	H_2O_2
15	ferryl iron	FeO^{2+}
	perferryl iron	FeO_4^{3-}


As used herein, "free radical damage" means the alteration in structure, function, composition, or other properties of biological tissues, organs, cells, or constituents that result from the effect of a free radical on them. Since free radicals are highly reactive and unstable species, they will in general react with a wide variety of biological targets to damage them. For example, the oxidation of lipids (lipid peroxidation), especially of cell membrane lipid, is a well-known damaging effect of radicals in biological systems.

As used herein, "controlling the onset of AIDS" means delaying, retarding, and/or preventing the activation of HIV types, especially HIV (1). The activation of HIV is often manifested by symptoms associated with AIDS. While not limited to any particular mechanism of action, the subject chelators are believed to control the onset of AIDS by reducing systemic levels of and/or preventing the formation of free radicals which play a role in the activation of the subject virus types.


Active Agent

The subject invention involves a method for protecting mammalian cells from free radical damage by reducing the level of free radicals in mammalian cells by administering to the mammal a safe and effective amount of an iron chelator selected from the group consisting of:


2-furildioxime having the structure:




2-furilmonoxime, having the structure:




1-phenyl-1,2-propanedione-2-oxime, having the structure:




5 1-phenyl-1,3-butanedione (benzoylacetone), having the structure:




1-hydroxy-4-methyl-6-(2,4,4-trimethylpentyl)-2(1H)-pyridinone (piroctone), having the structure:




10 diethyldithiocarbamic acid, having the structure:



deferoxamine, having the structure:



1,2-dimethyl-3-hydroxy-pyrid-4-one, (chelator L1), having the structure:



or a pharmaceutically-acceptable salt thereof; or mixtures of the subject compounds and/or their salts.

Preferred active compounds of the subject invention include 2-furildioxime, 2-furilmonoxime, 1-phenyl-1,2-propanedione-2-oxime (PPDO), piroctone and diethyl-dithiocarbamic acid. More preferred compounds include 2-furildioxime, 2-furilmonoxime, piroctone and diethyldithiocarbamic acid. Even more preferred compounds include 2-furildioxime and 2-furilmonoxime. The most preferred compound of the subject invention is 2-furildioxime.

Preferred actives useful in controlling the onset of AIDS include 2-furildioxime, 2-furilmonoxime, PPDO, benzoylacetone, piroctone, and 1,2-dimethyl-3-hydroxy-pyrid-4-one. More preferred actives useful in controlling the onset of AIDS include 2-furildioxime, 2-furilmonoxime, PPDO and piroctone. Even more preferred actives include 2-furildioxime and 2-furilmonoxime. The most preferred active is 2-furildioxime.

A particularly preferred piroctone salt is its ethanolamine salt, octopirox.

A particularly preferred salt of diethyldithiocarbamic acid is its sodium salt.

A particularly preferred salt of deferoxamine is its methanesulfonate salt, desferol.

Pharmaceutically-Acceptable Carrier

In addition to the active agent as described hereinbefore, the pharmaceutical compositions of the present invention essentially comprise a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier", as used herein, means one or more compatible solid or liquid filler diluents or encapsulating substances which are suitable for administration to a human or lower animal. The term "compatible", as used herein, means that the components of the pharmaceutical compositions are capable of being commingled with the compound of the present invention, and with each other, in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the pharmaceutical composition under ordinary use situations. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the human or lower animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers are sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives, such as sodium carboxymethylcellulose,

ethylcellulose, cellulose acetate; powdered tragacanth; malt; gelatin; talc; stearic acid; magnesium stearate; calcium sulfate; vegetable oils such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerine, sorbitol, mannitol, and polyethylene glycol; sugar; alginic acid; pyrogen-free
5 water; isotonic saline; phosphate buffer solutions; cocoa butter (suppository base); emulsifiers, such as the Tweens®; as well as other non-toxic compatible substances used in pharmaceutical formulations. Wetting agents and lubricants such as sodium lauryl sulfate, as well as coloring agents, flavoring agents, excipients, tableting agents, stabilizers, antioxidants, and preservatives, can also be present.

10 The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the compounds of the present invention is basically determined by the way the compound is to be administered. The preferred modes of administering the compounds of the present invention are orally, topically and by injection. Suitable pharmaceutically-acceptable carriers for topical application include those suited for use in creams, gels, tapes and the
15 like; and for oral administration include those suited for tablets and capsules.

The pharmaceutically-acceptable carrier employed in conjunction with the compounds of the present invention is used at a concentration sufficient to provide a practical size to dosage relationship. The pharmaceutically-acceptable carriers, in total, may comprise from about 0.1% to about 99.99% by weight of the pharmaceutical
20 compositions of the present invention, preferably from about 80% to about 99.9%, more preferably from about 90% to about 99.0%, even more preferably from about 92% to about 97%, also preferably from about 94% to about 96%.

Pharmaceutically-acceptable carriers suitable for the preparation of unit dosage forms for oral administration and injection, and dosage forms for topical application are
25 well-known in the art. Their selection will depend on secondary considerations like taste, cost, and/or shelf stability, which are not critical for the purposes of the subject invention, and can be made without difficulty by a person skilled in the art. Pharmaceutically-acceptable carriers useful in the compositions of the subject invention are described more fully hereinafter.

30 A. Oral Dose Forms:

Various oral dosage forms can be used, including such solid forms as tablets, capsules, granules, bulk powders and microcapsules of the drug. These oral forms comprise a safe and effective amount, usually at least about 5%, and preferably from about
35 10% to about 50% of the compound of the subject invention. Tablets can be compressed, enteric-coated, sugar-coated or filmcoated containing suitable binders, lubricants, surfactants, diluents, disintegrating agents, coloring agents, flavoring agents, preservatives, flow-inducing agents, and melting agents. Liquid oral dosage forms include aqueous and nonaqueous solutions, emulsions, suspensions, solutions and/or suspensions reconstituted

from non-effervescent granules, containing suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, melting agents, coloring agents, and flavoring agents. Preferred carriers for oral administration include gelatin and propylene glycol. Specific examples of pharmaceutically-acceptable carriers and excipients that may be used in formulating oral dosage forms containing compounds of the subject invention are described in U.S. Patent 3,903,297, Robert, issued September 2, 1975, incorporated by reference herein. Techniques and compositions for making solid oral dosage forms are described in Marshall, "Solid Oral Dosage Forms," Modern Pharmaceuticals, Vol. 7, (Banker and Rhodes, editors), 359-427 (1979), incorporated herein by reference. Techniques and compositions for making tablets (compressed, formulas and molded), capsules (hard and soft gelatin) and pills are described in Remington's Pharmaceutical Sciences (Arthur Osol, editor), 1553-1593 (1980), incorporated herein by reference.

The preferred unit dosage form for oral administration is tablets, capsules and the like, comprising a safe and effective amount of a compound of the subject invention. Preferably oral dose forms comprise from about 0.001 g to about 5g of a compound of the subject invention, more preferably from about 0.01 g to about 2g and most preferably from about 0.1g to about 1g, for a 50-70 kg person.

B. Topical Dose Forms:

The compositions of the subject invention can also be administered topically to a biological subject, i.e., by the direct laying on or spreading of the composition on the skin.

The topical compositions useful in the subject invention involve compositions suitable for topical application to mammalian skin, the composition comprising a safe and effective amount of an active agent or mixture of such actives as described hereinabove, and a pharmaceutically-acceptable topical carrier. The subject compositions preferably contain from about 0.001% to about 20%, preferably from about 0.01% to about 10%, more preferably from about 0.1% to about 5%.

The topical compositions useful in the subject invention may be made into a wide variety of product types. These include, but are not limited to lotions, creams, beach oils, gels, sticks; sprays, ointments, pastes, mousses, cosmetics, shampoos, cream rinses, hair tonics and hair conditioners. These product types may comprise several types of carrier systems including, but not limited to solutions, emulsions, gels, solids, and liposomes.

The topical compositions useful in the subject invention may include a safe and effective amount of penetration enhancing agent. A preferred amount of penetration enhancing agent is from about 1% to about 5% of the composition. Examples of useful penetration enhancers, among others, are disclosed in U.S. Patents 4,537,776, Cooper, issued August 27, 1985; 4,552,872, Cooper et al., issued November 12, 1985; 4,557,934, Cooper, issued December 10, 1985; 4,130,667, Smith, issued December 19, 1978; 3,989,816, Rhaadhyaksha, issued November 2, 1976; 4,017,641, DiGiulio, issued April

12, 1977; and 4,954,487, Cooper, Loomans & Wickett, issued September 4, 1990.

Other conventional skin care product additives may also be included in the compositions useful in the subject invention. For example, collagen, hyaluronic acid, elastin, hydrolysates, primrose oil, jojoba oil, epidermal growth factor, soybean saponins, mucopolysaccharides, and mixtures thereof may be used.

Various vitamins may also be included in the compositions useful in the subject invention. For example, Vitamin A, and derivatives thereof, Vitamin B₂, biotin, pantothenic, Vitamin D, and mixtures thereof may be used.

C. Injectable Dose Forms

The compounds of the subject invention are also useful when injected. The dosage of the compound of the subject invention which is both safe and effective will vary with the particular condition being treated, the severity of the condition, the duration of treatment, the specific compound employed and its usage concentration, and like factors within the specific knowledge and expertise of the attending physician and commensurate with a reasonable benefit/risk ratio associated with the use of any drug compound. The injectable dosages and dosage ranges given herein are based on delivery of the compound of the subject invention to a 70 kg human and can be adjusted to provide equivalent dosages for patients of different body weights.

Methods and materials for manufacturing injectables can be found in Remington's Pharmaceutical Sciences, 17ed., 1985, Chapter 85, p. 1518, the disclosures of which are incorporated herein by reference in their entirety. The injectable dosage forms typically contain from about 0.001 mg/ml to about 100 mg/ml, preferably from about 0.01 mg/ml to about 10 mg/ml, more preferably from about 0.1 mg/ml to about 3.0 mg/ml, of the compound of the subject invention. The injectable dosage forms are typically administered from about once a week to about four times daily, more preferably from about twice a week to about three times daily, more preferably still, about three times a week to about twice daily, also preferably about once daily. Typically, from about 1 ml to about 100 mls of the composition are injected, preferably from about 10 mls to about 50 mls, more preferably about 25 mls.

30 Combination Actives

A. Sunscreens and Sunblocks

Reduction of the level of free radicals in mammalian cells can be achieved by using combinations of the active agents together with sunscreens or sunblocks. A known inducer of free radicals is ultraviolet radiation. Thus, in topical compositions, the inclusion of sunscreens/sunblocks would increase protection against radical production and subsequent damage. Useful sunblocks include, for example, zinc oxide and titanium dioxide. The combination of an active agent, with a UVA and/or UVB sunscreen is desirable. The inclusion of sunscreens in compositions useful in the subject invention at

low levels does not greatly reduce the tanning response of the user but enhances the effectiveness of the subject compositions. A wide variety of conventional sunscreens agents are suitable for use in combination with a subject active agent. Sagarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and Technology, disclose numerous
5 suitable agents.

A safe and effective amount of sunscreen may be used in the compositions useful in the subject invention. The sunscreens agent must be compatible with the active agent. The composition preferably comprises from about 1% to about 20%, more preferably from about 2% to about 10% of a sunscreens agent. Exact amounts will vary
10 depending upon the sunscreen chosen and the desired Sun Protection Factor.

B. Anti-Inflammatory Agents

In a preferred composition useful in the subject invention, an anti-inflammatory agent is included as an active along with the active agent. The inclusion of an anti-inflammatory agent enhances the benefits of the compositions because the mammalian
15 body will respond to radical damage by mounting an inflammation response which can lead to additional cell damage. The anti-inflammatory agent also protects strongly in the UVA radiation range (though it also provides some UVB protection as well). (See U.S. Patent 4,847,071, Bissett, Bush, and Chatterjee, issued July 11, 1989, incorporated herein by reference; and U.S. Patent 4,847,069, Bissett and Chatterjee, issued July 11, 1989,
20 incorporated herein by reference.)

A safe and effective amount of an anti-inflammatory agent may be added to the compositions useful in the subject invention, preferably from about 0.1% to about 10%, more preferably from about 0.5% to about 5%, of the composition. The exact amount of anti-inflammatory agent to be used in the compositions will depend on the particular anti-inflammatory agent utilized since such agents vary widely in potency.
25

C. Anti-Oxidants/Radical Scavengers

In a preferred composition useful in the subject invention, an anti-oxidant/radical scavenger is included as an active along with a subject active agent. The inclusion of an anti-oxidant/ radical scavenger increases the benefits of the composition because an anti-oxidant/radical scavenger can neutralize any free radicals that are produced in spite of the subject actives' chelating action.
30

A safe and effective amount of an anti-oxidant/radical scavenger may be added to the compositions useful in the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition.

35 Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, tocopherol (vitamin E), tocopherol sorbate, other esters of tocopherol, butylated hydroxy benzoic acids and their salts, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (commercially available under the tradename Trolox®), gallic acid and its alkyl esters,

especially propyl gallate, uric acid and its salts and alkyl esters, sorbic acid and its salts, the ascorbyl esters of fatty acids, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine), sulfhydryl compounds (e.g., glutathione), and dihydroxy fumaric acid and its salts may be used.

- 5 In a preferred composition useful in the subject invention, compositions comprise one, any two, or all three of a sunscreensing agent, anti-inflammatory agent, and/or an anti-oxidant/radical scavenging agent included as actives along with a subject active agent. The inclusion of two or all three of these agents with the subject active increases the benefits of the composition.

10 D. Chelators

In a preferred composition useful in the subject invention, an additional chelating agent is included as an active along with a subject active agent. As used herein, "chelating agent" means an active agent capable of removing a metal ion from a system by forming a complex so that the metal ion cannot readily participate in or catalyze chemical reactions.

- 15 The inclusion of an additional chelating agent increases the benefits of the composition.

A safe and effective amount of a chelating agent may be added to the compositions useful in the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition. Chelators useful in compositions are disclosed in U.S. Patent Application Serial No. 619,805, Bissett, Bush & Chatterjee, filed November 27, 1990 (which is a continuation of U.S. Patent Application Serial No. 251,910, filed October 4, 1988); U.S. Patent Application Serial No. 514,892, Bush & Bissett, filed April 26, 1990; U.S. Patent Application Serial No. 657,847, Bush, Bissett & Chatterjee, filed February 25, 1991; and U.S. Patent Application Serial No. 776,506, Bush, filed October 11, 1991; all incorporated herein by reference. The additional
25 chelators preferred include:

- kojic acid
- 2,3-bis-(2-pyridyl)-pyrazine
- 3-(4-phenyl-2-pyridyl)-5-phenyl-1,2,4-triazine
- 2,3-bis-(2-pyridyl)-5,6-dihydropyrazine
- 30 2,4,6-tri-(2-pyridyl)-1,3,5-triazine
- 1-pyrrolidine carbodithioic acid
- di-2-pyridyl ketone
- phenyl 2-pyridyl ketoxime
- 2,3-dihydroxy naphthalene
- 35 2,3-dihydroxy pyridine
- 3-hydroxy-2-methyl-4-pyrone
- 2,3-dihydroxy benzoic acid
- ethylenediamine-N,N-bis-(2-hydroxy-phenylacetic acid) dimethyl ester

In a preferred composition useful in the subject invention, compositions comprise one, any two, any three, or all four of a sunscreensing agent, anti-inflammatory agent, anti-oxidant/radical scavenging agent, and/or chelating agent included as actives along with the subject active agent. The inclusion of two, three, or all four of these agents with the active agent increases the benefits of the composition.

Methods for Reducing Free Radical Damage in Mammalian Cells

The subject invention relates to methods for reducing the level of free radicals in mammalian cells. The reduction of the level of free radicals in mammalian cells reduces the level of free radical aging in the cells. Reduced levels of free radicals also prolongs the interval between infection by HIV in certain mammals, and the activation of the HIV (and the accompanying symptoms of AIDS exhibited by the HIV positive mammal).

The subject methods comprise topical application of a safe and effective amount of an active agent. The amount of treatment and frequency of application will vary widely depending upon the conditions of the cells already in existence in the subject and the level of treatment desired.

A safe and effective amount of active agent, in a topical composition, is applied, generally from about 0.001 mg to about 2 mg per cm² skin per application, preferably from about .005 mg to about 1 mg per cm² skin per application, more preferably from about 0.01 mg to about 0.5 mg/cm², also preferably from about 0.02 mg to about 0.2 mg/cm². Application preferably ranges from about weekly to about 5 times daily, more preferably from about twice a week to about four times daily, more preferably still from about every other day to about 3 times daily, also preferably from about once a day to about twice a day. The subject compositions are preferably applied to an area of from about 10 cm² to about 10,000 cm² skin for each application, more preferably from about 100 cm² to about 5,000 cm² skin, also preferably from about 500 cm² to about 1000 cm² skin. Treatment is continued for at least 7 days, more preferably 6 months, even more preferably 1 year, more preferably still 5 years, also preferably 10 years.

A preferred method of the subject invention involves applying both a safe and effective amount of the active and a safe and effective amount of one or more of a sunscreensing agent, an anti-inflammatory agent, an anti-oxidant/radical scavenging agent, and/or a chelating agent, to the skin simultaneously. As used herein, "simultaneous application" or "simultaneously" means applying the agents to the skin at the same situs on the body at about the same time. Though this can be accomplished by applying the agents separately to the skin, preferably a composition comprising all the desired agents commingled is applied to the skin. The amount of sunscreensing agent applied is preferably from about 0.05 mg to about 0.5 mg per cm² skin. The amount of anti-inflammatory agent applied is preferably from about 0.005 mg to about 0.5 mg, more preferably from about 0.01 mg to about 0.1 mg per cm² skin. The amount of anti-oxidant/radical

scavenging agent preferably applied is from about 0.01 mg to about 1.0 mg, more preferably from about 0.05 mg to about 0.5 mg per cm² skin. The amount of chelating agent preferably applied is from about .001 mg to about 1.0 mg, more preferably from about 0.01 mg to about 0.5 mg, still more preferably from about 0.05 mg to about 0.1 mg per cm² skin. The amount of the active compound of the subject invention applied is
5 preferably from about 0.001 mg to about 2 mg per cm² skin per application, more preferably from about 0.01 mg to about 1 mg per cm² skin per application.

The preferred modes of administration are orally, topically, and parenterally (for example, by subcutaneous injection, intramuscular injection, intra-articular injection,
10 intravenous injection and the like). Thus, specific modes of administration include, without limitation, oral, transdermal, mucosal, sublingual, intramuscular, intravenous, intraperitoneal, and subcutaneous administration, as well as topical application.

Oral administration can be used to reduce the level of free radicals through oral dosing of a pharmaceutical composition comprised of a safe and effective amount of the
15 compound of the subject invention in a suitable oral pharmaceutical carrier. The compound is absorbed by the gastrointestinal tract. The pharmaceutical composition may consist of solid dosage forms such as tablets, hard gelatin capsules, soft gelatin capsules, bulk powders, and microcapsules of the drug. Alternately, it may consist of a liquid dosage form such as an aqueous or nonaqueous solution, emulsion, or suspension.

The amount of the compound ingested depends upon the bioavailability of the compound from the oral pharmaceutical composition. Typically, however, the compounds of the subject invention are dosed in an amount of from about 0.001 mg/kg of body weight to about 100 mg/kg, preferably from about 0.01 to about 50 mg/kg of body weight, more preferably from about 0.1 to about 20 mg/kg of body weight, also preferably from about 1
25 mg to about 10 mg/kg of body weight. The amount of the pharmaceutical composition depends upon the percent of compound within its formula, which is a function of the amount of the compound required per dose, its stability, release characteristics and other pharmaceutical parameters. Generally, the oral pharmaceutical composition should comprise from about 5% to about 50% of the compound of the subject invention.

Oral application preferably ranges from about weekly to about 5 times daily, more preferably from about twice a week to about four times daily, more preferably still from about every other day to about 3 times daily, also preferably from about once a day to about twice a day. Treatment is continued for at least 7 days, more preferably 6 months,
30 even more preferably 1 year, more preferably still 5 years, also preferably 10 years.

For controlling the onset of AIDS, the compounds of the subject invention are preferably dosed in an oral form in an amount of from about 10 mg to about 1000 mg per m² of body surface area of the subject infected with HIV, more preferably from about 100 to about 800 mg per m², also preferably from about 300 to about 500 mg per m², most
35

preferably about 400 mg per m². Application preferably ranges from about once a month to about twice daily, more preferably about once every two weeks to about once a day, more preferably still about once every 10 days to about once every other day, also preferably about once a week.

- 5 The preferred method of injectable administration depends upon the solubility and the stability of the particular active being used.

EXAMPLES

- 10 The following examples further describe and demonstrate embodiments within the scope of the subject invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the subject invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

ORAL DOSE FORMS

EXAMPLE I

- 15 A composition for oral administration is prepared by combining the following

Component:

2-furildioxime (FDO)	1 kg
Sesame oil	to 4 liters

- 20 The FDO is suspended in the sesame oil with the aid of sonication and is packaged in soft gelatin capsules using methods known in the art. Two of the resulting capsules, each containing 250 mg of the active, are administered to a 60 kg human, daily for a period of 4 years.

EXAMPLE II

- A composition for oral administration is prepared by combining the following :

Component

25	2-furilmonoxime (FMO)	250 g
	Propylene glycol	1800 ml
	Ethyl alcohol	175 ml
	Distilled water	75 ml
	Artificial Cherry flavor	10 ml
30	FD&C Red #40	0.2 g

- The above ingredients are combined to produce a syrup and are packaged under sterile conditions in 6 oz. bottles. One teaspoon of this formulation is administered to a 70 kg human, weekly for a period of one year.

EXAMPLE III

- 35 Tablets are prepared by conventional methods, such as mixing and direct

compaction, formulated as follows:

<u>Component</u>	<u>mg per tablet</u>
piroctone	500
Microcrystalline cellulose	400
5 Sodium Starch glycolate	60
Magnesium stearates	10

One tablet is administered orally to a human infected with HIV, once a week for a period of one year or more.

TOPICAL DOSE FORMS

10

EXAMPLE IV

A topical composition is prepared by combining the following components utilizing conventional mixing techniques.

<u>Component</u>	<u>% Weight</u>
Ethanol:propylene glycol:water (1:1:2)	98.7
15 Diethyldithiocarbamic acid(DEDCA), sodium salt	1.3

In a suitable vessel, the Na-DEDCA is dissolved in the ethanol:propylene glycol:water with stirring. Use of an amount of the composition to deposit about 0.02 mg/cm² of the active agent to about 200 cm² of skin is appropriate. The composition is
20 applied twice daily, for a period of five years.

EXAMPLE V

A topical composition is prepared by combining the following components utilizing conventional mixing techniques.

<u>Component</u>	<u>% Weight</u>
25 Ethanol	99.87
PPDO	0.13

In a suitable vessel, the PPDO is dissolved in the ethanol with stirring. Use of an amount of the composition to deposit about 0.001 mg/cm² of the active agent to the skin is appropriate. The composition is applied four times daily over a 1000 cm² area of skin
30 for a period of six months.

EXAMPLE VI

A topical composition is prepared by combining the following components utilizing conventional mixing techniques.

<u>Component</u>	<u>% Weight</u>
35 Ethanol	49.00

Propylene glycol	25.00
Deionized water	25.00
Octopirox	1.00

In a suitable vessel, the octopirox is dissolved in the ethanol with stirring. Propylene glycol and deionized water are added with stirring. Use of an amount of the composition to deposit about 0.02 mg/cm^2 of the active agent to 2000 cm^2 of skin is appropriate. The composition is applied once a week for one year.

EXAMPLE VII

A nonionic oil-in-water emulsion is prepared by combining the following components utilizing conventional mixing techniques:

	<u>Component</u>	<u>% Weight</u>
	Deionized Water	79.73
	Propylene Glycol	3.00
	Octyl Methoxycinnamate	7.50
15	Cetyl Alcohol	2.50
	Stearyl Alcohol	2.50
	Laureth 23	2.00
	C ₁₂₋₁₅ Alcohols Benzoate	2.00
	EDTA	0.37
20	Methylparaben	0.20
	Propylparaben	0.10
	Deferoxamine	0.10

Use of an amount of the composition sufficient to deposit about 0.004 mg/cm^2 skin of the active agent is appropriate. The composition is applied twice a week to about 500 cm^2 of skin for two years.

EXAMPLE VIII

A nonionic oil-in-water emulsion is prepared by combining the following components utilizing conventional mixing techniques:

	<u>Component</u>	<u>% Weight</u>
30	Deionized Water	78.73
	Propylene Glycol	3.00
	Octyl Methoxycinnamate	7.50
	Cetyl Alcohol	2.50
35	Stearyl Alcohol	2.50
	Laureth 23	2.00
	C ₁₂₋₁₅ Alcohols Benzoate	2.00

EDTA	0.37
Methylparaben	0.20
Propylparaben	0.10
Benzoylacetone	1.10

- 5 Use of an amount of the composition sufficient to deposit about 0.05 mg/cm² of the active agent to about 100 cm² of skin is appropriate. The composition is applied once daily, for three years.

EXAMPLE IX

- 10 An ion pair oil-in-water emulsion is prepared by combining the following components utilizing conventional mixing techniques.

<u>Component</u>	<u>% Weight</u>
Deionized Water	78.05
Permuron TR-2 (C10-C30 Acrylate Copolymer, B.F. Goodrich)	0.30
15 Distearyl Dimethyl Ammonium Chloride	0.15
1,2-dimethyl-3-hydroxy-pyrid-4-one	1.00
4-N,N-(2-ethylhexyl)methylaminobenzoic acid 4-ester of 2-hydroxy-4-(2-hydroxyethoxy)- benzophenone	4.00
20 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane	2.00
Dimethyl Isosorbide	6.00
Dioctyl Malate	6.00
Cetyl Alcohol	1.00
25 Stearyl Alcohol	1.00
99% Triethanolamine	0.50

- Use of an amount of the composition sufficient to deposit about 0.1 mg/cm² of the active agent to about 1000 cm² of skin is appropriate. The composition is applied twice daily, for one year.

30

EXAMPLE X

- A sunscreen composition is prepared by combining the following components utilizing conventional mixing techniques.

<u>Component</u>	<u>% Weight</u>
Polypropylene Glycol 15 Stearyl Ether	15.00
35 Sorbitan Oleate	2.00
Octyl Methoxy Cinnamate	7.50
FDO	0.50

	Propyl Paraben	0.15
	Butylated Hydroxy Toluene	0.05
	Cyclomethicone	20.00
	Sesame Oil	5.00
5	Mineral Oil (Blandol)	49.80

Use of an amount of the composition is sufficient to deposit about 0.1 mg/cm² of the active agent to the skin is appropriate. This composition is applied three times daily, for six months.

10 While particular embodiments of the subject invention have been described, it will be obvious to those skilled in the art that various changes and modifications of the subject invention can be made without departing from the spirit and scope of the invention. It is intended to cover, in the appended claims, all such modifications that are within the scope of the invention.

What is claimed is:

1. A method of reducing free radical damage in cells of living mammals comprising administering to a mammal a composition comprising a safe and effective amount of an iron chelator selected from the group consisting of 2-furildioxime, 2-furilmonoxime, 1-phenyl-1,2-propanedione-2-oxime, benzoylacetone, piroctone, diethyldithiocarbamic acid, deferoxamine and 1,2-dimethyl-3-hydroxy-pyrid-4-one; or pharmaceutically-acceptable salts thereof, preferably selected from the group consisting of 2-furildioxime, 2-furilmonoxime, 1-phenyl-1,2-propanedione-2-oxime, piroctone and diethyldithiocarbamic acid, more preferably selected from the group consisting of 2-furildioxime, 2-furilmonoxime, piroctone and diethyldithiocarbamic acid, and most preferably wherein the iron chelator is 2-furildioxime or 2-furilmonoxime.
2. The method of any of Claims 1 wherein the composition is administered in an oral dose form wherein from 0.001 mg to 100 mg of the chelator, preferably from 0.01 mg to 50 mg most preferably from 0.1 mg to 20 mg is administered per kg of body weight of a human, from once a week to 5 times daily, for a period of one week or more, preferably from twice a week to four times daily, for a period of one year or more, most preferably from once a day to twice a day, for a period of five years or more, preferably 10 years or more.
3. The method of any of Claims 1 wherein the composition is applied topically to the skin of the mammal wherein the amount of chelator applied to the skin is from .001 mg to 2 mg per cm² skin, preferably from 0.005 mg to 1 mg per cm² skin most preferably from 0.01 mg to .5 mg per cm² skin; to an area of 10 cm² to 10,000 cm² of skin, preferably to an area of 100 cm² to 5000 cm² skin, most preferably to an area of 500 cm² to 1000 cm² skin for each application, from once a week to 5 times daily, for a period of one week or more, preferably from twice a week to 4 times daily, for a period of one year or more, most preferably from once a day to twice a day, for a period of five years or more, preferably 10 years or more.
4. A method of reducing the level of free radicals in mammalian cells comprising administering to a mammal a composition comprising a safe and effective amount of an iron chelator selected from the group consisting of 2-furildioxime, 2-furilmonoxime, 1-phenyl-1,2-propanedione-2-oxime, benzoylacetone, piroctone, diethyldithiocarbamic acid, deferoxamine and 1,2-dimethyl-3-hydroxy-pyrid-4-one; or a pharmaceutically-acceptable salt thereof.
5. A method of controlling the onset of AIDS in a mammal infected by HIV comprising administering to the mammal a safe and effective amount of an iron chelator selected

from the group consisting of 2-furildioxime, 2-furilmonoxime, 1-phenyl-1,2-propanedione-2-oxime, benzoylacetone, piroctone, and 1,2-dimethyl-3-hydroxy-pyrid-4-one; or a pharmaceutically-acceptable salt thereof.

6. The method of Claim 5 wherein the chelator is administered in an oral dosage form, in an amount of from 10 mg to 1000 mg per m², preferably from 100 mg to 800 mg per m², most preferably from 300 mg to 500 mg per m² body surface area of the mammal, from once a month to twice daily, for a period of one month or more, preferably from once every two weeks to once daily, for a period of five years or more, most preferably from once every ten days to once every other day, for a period of ten years or more.
7. The method of any of Claims 1 through 6 wherein the chelator administered is 2-furildioxime.

INTERNATIONAL SEARCH REPORT

International application No. . . .
PCT/US95/02872

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K, 31/335, 31/34, 31/27, 31/13, 31/12, 31/16

US CL : 514/350, 471, 476, 616, 641, 678

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/350, 471, 476, 616, 641, 678

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
STN-compound; Antiviral use & Free Radicul degree use.Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
as above.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, VOLUME 115, NO. 4, ISSUED 12 MARCH 1991, HALLIWELL, "FREE RADICAL BIOL. MED", 1989, 7(6), 645-651, COLUMN 1, ABSTRACT NO. 21476t, SEE ENTIRE DOCUMENT.	1-4
Y	CHEMICAL ABSTRACT, VOLUME 118, NO. 12, ISSUED 26 JULY 1993, FEORINO ET AL. "ANTIVIRAL CHEM. CHEMOTHER.", 1993, 4(1), 55-63, COLUMN 2, ABSTRACT NO. 182829 n, SEE ENTIRE DOCUMENT.	5-6

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A		document defining the general state of the art which is not considered to be of particular relevance
* E		earlier document published on or after the international filing date
* L		document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
* O		document referring to an oral disclosure, use, exhibition or other means
* P		document published prior to the international filing date but later than the priority date claimed
	* X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
	* Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
	* &	document member of the same patent family

Date of the actual completion of the international search 05 JUNE 1995	Date of mailing of the international search report 22 JUN 1995
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer RUSSELL TRAVERS aco Telephone No. (703) 308-1235